

American
Urological
Association

Education and Research, Inc.

Practice Guidelines from the American Urological Association

AUA GUIDELINES

A Department within the Health Policy Division



American
Urological
Association

Education and Research, Inc.

The American Urological Association

- **The American Urological Association (AUA), founded in 1902, is the premier professional association for the advancement of urologic patient care, and works to ensure that its more than 18,000 members are current on the latest research and practices in urology.**
- **AUA Mission Statement:**
 - **To promote the highest standards of urological clinical care through education, research and in the formulation of health care policy.**

AUA GUIDELINES

A Department within the Health Policy Division



American
Urological
Association

Education and Research, Inc.

AUA Guidelines

- **Health Policy/Science and Quality Division**
- **Practice Guidelines Committee**
- **Six Staff Positions**

AUA GUIDELINES

A Department within the Health Policy Division



American
Urological
Association

Education and Research, Inc.

AUA Guidelines: Current Process

- 1. Topic Nomination**
- 2. Panel Identification**
- 3. Definition of Research Questions**
- 4. Literature Review & Appraisal**
- 5. Data Synthesis & Analysis**
- 6. Evidence Report & Guideline Statement Development**
- 7. Writing**
- 8. Peer Review & Revision**
- 9. Approval & Publication**

AUA GUIDELINES

A Department within the Health Policy Division



American
Urological
Association


Education and Research, Inc.

AUA Guidelines: Dissemination:

- **Dissemination for all new guidelines:**
 - Online publication
 - NGC/GIN submission
 - Plenary presentation at Annual Meeting
 - Executive Summary for Journal of Urology
 - Guidelines-at-a-glance (print and app)
 - Primary Care Pocket Guides
 - Translated Guidelines
 - Pocket table/Wall chart
 - Webinar
 - AUA News/HPBrief article
 - Press release
 - Social media announcements
 - Email blast
 - Slide set



Connect with the AUA



AUA GUIDELINES

A Department within the Health Policy Division



American
Urological
Association

Education and Research, Inc.

AUA Guidelines: Dissemination

- **Smartphone Applications**
- **Video Panel Discussions**
- **Adaptive Spaced Education (Qstream)**




AUA GUIDELINES

A Department within the Health Policy Division

AUA Guidelines: EHR integration

- **GLIDES workgroup**
 - **BRIDGE-Wiz**
 - **eGLIA**
 - **GEM cutter**
- **Panel Informaticists**
 - **XML formatting**
- **EHR vendors**





American
Urological
Association

Education and Research, Inc.

BRIDGE-Wiz



- Formalizes a process for writing implementable recommendations
- Focuses discussion
- Incorporates prompts based on COGS to improve guideline quality
- Controlled natural language
 - Offers verb choices based on action-type
 - Traps and disallows use of “consider”
 - Discourages “statement of fact” masquerading as recommendation
 - Limits boolean connectors to all ANDs or ORs in a statement
- Incorporates decidability and executability checks
- Requires systematic appraisal of evidence quality and benefit-harms
 - Suggests appropriate obligation term (deontic modal)
- Output includes a high-level “rule” and an evidence profile

AUA GUIDELINES

A Department within the Health Policy Division



Add Another Action

What

What type of activity do you propose?

EDUCATE/COUNSEL

NOT

Based on the EDUCATE/COUNSEL action type,
select a verb:

recommend

ADD

Recommend what?

diet and exercise

VERB

WHAT

	start	metformin as first-line treatment	AND	OR	DEL
AND	recommend	diet and exercise	AND	OR	DEL



American
Urological
Association

Education and Research, Inc.

Define the “Actor”

Who

WHO is to start metformin as first-line treatment AND recommend diet and exercise IF newly diagnosed with type 2 diabetes, i.e., not previously treated?

Clinicians



Select Aggregate Evidence Quality

Evidence Quality

What aggregate evidence quality supports these benefits, risks, harms, and costs?
start metformin as first-line treatment AND recommend diet and exercise

IF
newly diagnosed with type 2 diabetes, i.e., not previously treated

Evidence Quality

- A Well-designed, well-conducted randomized, controlled trials or diagnostic studies performed on a population similar to the guideline's target population
- B Randomized, controlled trials with "nonfatal flaws" or methodologic limitations; overwhelmingly consistent evidence from observational studies
- C Observational studies (case control or cohort design)
- D Expert opinion, case reports, reasoning from first principles
- X Exceptional situations where validating studies cannot be performed and there is a clear preponderance of harm or benefit

Resources

Quality Appraisal

Aggregate

RCT-Consort

RCT-Casp

Observation

MetaAnalysis

Diagnostic

Notes

Back

Next

Restart

Judge Benefit-Harms Balance

Balance

Is there Equilibrium or a preponderance of Benefits or Risks, Harms, and Costs?



- Equilibrium
- Preponderance of Risks, Harms, Costs
- Preponderance of Benefit

Benefits

Lower Hgb A1c
 Target A1c sustained longer
 Less early deterioration of blood glucose
 Lower chance of weight gain
 Improved insulin

SMBG

Possible metabolic deterioration if Type 1 is misdiagnosed and treated as type 2
 Potential risk of lactic acidosis in setting of ketosis or significant dehydration



American
Urological
Association

Education and

Choose a Recommendation Style

If
newly-diagnosed with type 2 diabetes (i.e., not previously treated)
Then
Clinicians should start metformin as first-line treatment AND
recommend diet and exercise

Select

Clinicians should start metformin as first-line treatment AND
recommend diet and exercise
if/when/whenever
newly-diagnosed with type 2 diabetes (i.e., not previously treated)

Select

The *{developer}* recommends that if
newly-diagnosed with type 2 diabetes (i.e., not previously treated)
Then
Clinicians should start metformin as first-line treatment AND
recommend diet and exercise

Select

The *{developer}* recommends that
Clinicians should start metformin as first-line treatment AND
recommend diet and exercise
if/when/whenever
newly-diagnosed with type 2 diabetes (i.e., not previously treated)

Select



American
Urological
Association

Education and Research, Inc.

Urodynamics Panel Experience

- **Orientation at panel meeting (May 2011)**
- **Learning curve: Practice makes perfect**
- **Visual representation of benefits vs risks/burden was helpful**
- **Adaptable to AUA system**



Results

- **Clinicians who are making the diagnosis of urodynamic stress incontinence should assess urethral function.
(*Recommendation; Evidence Strength: Grade C*)**
- **Clinicians should perform repeat stress testing with the urethral catheter removed in patients suspected of having SUI who do not demonstrate this finding with the catheter in place during urodynamic testing.
(*Recommendation; Evidence Strength: Grade C*)**

Comparison: Overactive Bladder Guideline

- **In rare cases, augmentation cystoplasty or urinary diversion for severe, refractory, complicated OAB patients may be considered. *Expert Opinion***
- **Behavioral therapies may be combined with anti-muscarinic therapies. *Recommendation***

Next Steps

- **eGLIA: evaluation of Implementability**
- **GEM cutter III: XML format**
- **Clinical Decision Support**



American
Urological
Association

Education and Research, Inc.

- **Questions?**
- **Comments?**

AUA GUIDELINES

A Department within the Health Policy Division